

Comparative anatomy of adductors thigh muscles in capuchin monkey

Paula, JP., Ferreira, TAA., Paula, JP., Silva, MSL., Prado, YCL. and Marin, KA.

Universidade Federal de Goiás

The behavioral comparative analysis of recent primates, associated with the phylogenetic trunks derivative from cladism, generate important knowledge of cognitive evolution human recent, mainly associated with no humans primates from Old World, however, the recent studies on learning and memory of the *Cebus*, demonstrated that this animal is put joint with these others Old World primates, because *Cebus* and Old World primates have the same basic neural substrate for memory and learning tests indicating a long time convergent development of these species. The comparative anatomical studies between the *Cebus* and chimpanzees done by us corroborates these data of evolutionary convergence concerned to the muscles, nerves and vessels of thoracic members. The mainly evolutionary characters acquired by hominids were the erect posture and bipedal locomotion on the soil substituting the brachiation (hand used to locomotion on the trees). There are evolutionary and morphologic similarities between *Cebus* and humans and big similarities in bipedal nature. The survive and arboreous behavioral of no humans primates, indicate morphological similarities between these species and *Cebus*. The anatomical study provides substrate for verification of motors abilities of primates' species considering the muscles numbers, division and individualization of muscular body that insert in osseous portion. The purpose of this work is study the thigh muscles of *Cebus* genus and to compare the results with literature data of these muscles in humans, chimpanzees and baboons, and associate these results with any behavioral aspects. These data will provide the morphological differences and similarities between the *Cebus*, Old World primates and humans, contributing to correlations possible on evolutionary, phylogenetic and behavioral aspects between these species. In this study, 8 *Cebus libidinosus* monkeys were used, all healthy adults, with deviations as for size and age. They were provided by IBAMA (Brazilian Institute for the Environment and Renewable Natural Resources), from the city of Sete Lagoas, Minas Gerais State in 1970, and housed at the anatomy collection of the Federal University of Goiás (UFG). All animals were prepped with injections of Neoprene 601A latex (DuPont) on the femoral artery. The animals were carefully dissected using the naked eye or with the aid of 10x stereoscopic magnification. The muscles located on medial side of thigh, all with adductor action were 1) pectineus, 2) adductor longus, 3) adductor magnus, 4) adductor brevis, and gracile. The adductor brevis, adductor longus and gracile in *Cebus* have different origins in studied, but the insertions are identical to chimpanzees, baboons and *Cebus*, but not to human. The general aspects of these muscles to *Cebus* are more similar to baboons, putatively to arboreous behavioral similar between these last monkeys.

Financial support: FUNAPE, CNPq, CAPES, IACT.